



## **UWL REPOSITORY**

**[repository.uwl.ac.uk](https://repository.uwl.ac.uk)**

Applying the common-sense model to irritable bowel syndrome: the role of illness and treatment representations

Usher, Lee, Fox, Pauline ORCID: <https://orcid.org/0000-0003-0046-4940> and Mitchell, Kathryn (2012) Applying the common-sense model to irritable bowel syndrome: the role of illness and treatment representations. In: European Health Psychology Society Conference 2012, 21-25 Aug 2012, Prague, Czech Republic.

<http://dx.doi.org/10.1080/08870446.2012.707817>

**This is the Accepted Version of the final output.**

**UWL repository link:** <https://repository.uwl.ac.uk/id/eprint/1897/>

**Alternative formats:** If you require this document in an alternative format, please contact: [open.research@uwl.ac.uk](mailto:open.research@uwl.ac.uk)

### **Copyright:**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy:** If you believe that this document breaches copyright, please contact us at [open.research@uwl.ac.uk](mailto:open.research@uwl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

Title: Applying the common-sense model to Irritable Bowel Syndrome: the role of illness and treatment representations.

Lee Usher, Pauline Fox and Kathryn Mitchell. University of West London.

Background: Theorised pathways in an 'extended' common-sense model of illness representations (CSM) in those affected by Irritable Bowel Syndrome (IBS) were examined. Analysis contrasted between those using complementary medicines (CAM-users) and those not using CAM (non-users).

Methods: A web-based cross-sectional survey. Participants (n=653) were recruited from an IBS self-help network and other online illness discussion forums and assessed on the IPQ-R, the BMQ-General, the Brief-COPE and IBS-QOL.

Findings: CAM-users reported poorer quality of life compared to non-users. Stronger perceptions of illness consequences and emotional representations were strongly correlated with reduced quality of life and poorer emotional outcomes in both groups. Self-blame and behavioural disengagement partially mediated the pathway from perception to outcome.

Discussion: Findings suggest consistency with CSM theory. The detected influence of representations on quality of life (via coping strategies) suggests health psychologists could address components of illness perceptions to improve quality of life in those affected by IBS.